

# Pulsational frequencies of the eclipsing $\delta$ Scuti star HD 172189. Results of the STEPHI XIII campaign.

J. E. S. Costa<sup>1</sup>, E. Michel<sup>1</sup>, J. Peña<sup>2</sup>, O. Creevey<sup>3</sup>, Z. P. Li<sup>4</sup>, M. Chevreton<sup>1</sup>, J. A. Belmonte<sup>3</sup>, M. Alvarez<sup>2</sup>, L. Fox Machado<sup>3</sup>, L. Parrao<sup>2</sup>, F. Pérez Hernández<sup>3</sup>, A. Fernández<sup>1</sup>, J. R. Fremy<sup>1</sup>, S. Pau<sup>1</sup>, and R. Alonso<sup>3</sup>

<sup>1</sup> Observatoire de Paris, LESIA, FRE 2461, 92195 Meudon, France

<sup>2</sup> Instituto de Astronomía - Universidad Nacional Autónoma de México, Ap.P. 877, Ensenada, BC, Mexico

<sup>3</sup> Instituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain

<sup>4</sup> National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

Received 00.00.0000 / Accepted 22.02.2007

## ABSTRACT

**Context.** The eclipsing  $\delta$  Scuti star HD 172189 is a probable member of the open cluster IC 4756 and a promising candidate target for the CoRoT mission.

**Aims.** The detection of pulsation modes is the first step in the asteroseismological study of the star. Further, the calculation of the orbital parameters of the binary system allows us to make a dynamical determination of the mass of the star, which works as an important constraint to test and calibrate the asteroseismological models.

**Methods.** We performed a detailed frequency analysis of 210 hours of photometric data of HD 172189 obtained from the STEPHI XIII campaign\*.

**Results.** We have identified six pulsation frequencies with a confidence level of 99% and a seventh with a 65% confidence level, in the range between 100 – 300  $\mu$ Hz. In addition, three eclipses were observed during the campaign, allowing us to improve the determination of the orbital period of the system.

**Key words.** Stars: oscillations (including pulsations) — (Stars: variables:)  $\delta$  Sct — (Stars:) binaries: eclipsing

## 1. Introduction

HD 172189 (IC 4756 93 = SAO 123754) is a binary star of visual magnitude  $m_V = 8.85$  and spectral type A2 ( $\alpha_{2000} = 18^h38^m37.554^s$ ,  $\delta_{2000} = +05^\circ27'55.34''$ ) in the galactic cluster IC 4756. Recently, Martín-Ruiz et al. (2005) (M-R, hereafter) showed that HD 172189 is an eclipsing binary star with an orbital period of 5.71 days and with  $\delta$  Scuti-type pulsations with a clear frequency of 19.5974  $\text{cd}^{-1}$  (226.8  $\mu$ Hz), and evidence of more modes in the range 208 – 232  $\mu$ Hz.

The  $\delta$  Scuti variables are stars with masses from 1.5 to 2.5  $M_\odot$ , located at the intersection of the lower part of the classical Cepheid instability strip with the main sequence. They consist of main sequence objects, pre-main sequence ones, and also objects that have evolved off the main sequence, in the hydrogen shell burning phase. Most of the  $\delta$  Scuti stars are multiperiodic, with frequencies between 50 and 600  $\mu$ Hz. The pulsation modes are radial as well as non-radial due to  $\kappa$ -mechanism associated with the zone where He is partially ionized. Thus,  $\delta$  Scuti stars provide a good opportunity to apply asteroseismology and study key mechanisms at work in the main sequence stage, among which are transport of angular momentum, transport of chemical species at the edge of convective cores (overshooting), and large-scale circulations.

A few weeks of observations are usually required to resolve their rich spectra, often showing close pairs of peaks with a

spacing of  $\sim 0.5 \mu\text{Hz}$ . Three week-long multisite observational campaigns are regularly organized within the STEPHI network (Michel et al. 1992; Michel et al. 2000), trying to minimize the problem of missing data due to the day-night cycle, which induces strong side lobes of the spectral window in the Fourier spectrum.

The frequency analysis of HD 172189 shows that it is a  $\delta$  Scuti with at least six pulsation frequencies. Eclipses that occurred during the campaign were observed, allowing us to refine the period of the binary system.

The star HD 172189 was found to be a "likely member" of the open cluster IC 4756 by Kopff (1943). Herzog et al. (1975) updated the results of a previous work of Seggewiss (1968), concluded that the probability of the membership to the cluster for HD 172189, based on its proper motion alone, is 92%. Missana & Missana (1995) re-studied the cluster IC 4756, estimating the probability of membership of each stars and arrived to a similar result for the case of HD 172189: 95% of probability of being a member of the cluster.

The interest of pulsating stars in cluster and/or in well characterized binaries for asteroseismology studies has been stressed for long and by several authors. As mentioned in a recent review by Aerts (2006) (see also Aerts and Harmanec 2004), beside the perspective to study processes specific to binarity, like the influence of tidal forces on angular momentum transport or eventual mass transfer, the possibility to determine precisely global parameters (like individual masses, radii, chemical composition or age) allows to reduce significantly the parameters space to explore in the necessary stellar modeling work. The seismic inter-

Send offprint requests to: J.E.S. Costa

\* The HD 172189 reduced light curves are available in the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>.

pretation is thus expected to become more constraining in terms of diagnostics on physical processes.

In this spirit, Brown et al. (1994) applied singular value deconvolution technique to study the relative impact of precision on eigenfrequencies and global parameters in the illustrative case of two solar-like pulsators in a visual binary. They considered the optimal case of the close Alpha Cen binary. Their conclusions necessarily remained theoretical, since no oscillations were detected at that time neither in Alpha Cen A nor in Alpha Cen B. But this seminal work found a great development in Miglio and Montalbán (2005) where an optimization using Levenberg-Marquardt algorithm is applied to Alpha Cen A+B, considering all available observables, including oscillation frequencies now detected in both objects. This work confirms that satisfying all the observables with present physical assumptions is becoming more and more challenging and ineluctably leads to question the description of key-physical processes like mixing length theory versus full spectrum theory. Miglio and Montalbán however conclude that the present seismic data do not allow to go further and discriminate between different physical options for the equation of state or element diffusion descriptions.

In the specific case of  $\delta$  Scuti stars, Creevey et al. (2006) proposed to develop the approach of Brown et al. (1994), in the specific case of eclipsing binaries featuring a  $\delta$  Scuti. Their work is however, so far limited only to non seismic observables.

A few  $\delta$  Scuti pulsators in eclipsing binaries have already been intensively searched for oscillations. For RZ Cas and AB Cas (Rodríguez et al. 2004a; Rodríguez et al. 2004b), these observations only brought one oscillation frequency, and they did not lead to a seismic interpretation so far. RS Cha is another example of eclipsing binary featuring pre-main sequence objects, one of them showing  $\delta$  Scuti type oscillations (Alecian et al. 2006). Recently, new observations have been organized successfully, showing apparent multiperiodicity of both component (Bohm et al. In prep.).

The fact that HD 172189 is a  $\delta$  Scuti star and at the same time a binary system and a member of the open cluster IC 4756 makes it a very interesting target for the CoRoT mission.

## 2. The observations

The STEPHI campaigns include three observatories around the world: Observatorio del Teide (Izaña, Tenerife Island, Spain), Observatorio Astronómico Nacional de San Pedro Mártir (Baja California, Mexico) and the Xinglong Station of the Beijing Observatory (China). The observations were carried out for 21 days over the period 7-28 June 2004. The B9 star HD 172248 (IC 4756 117 = SAO 123762) was used as the primary comparison star (comp1) with a brightness of  $m_V = 8.91$ . Two secondary comparison stars (comp2) were observed: IC 4756 142 at the Mexican observatory and IC 4756 115 at the Spanish observatory. The first one is an A3 star with  $m_V = 9.51$  while the latter is a star of spectral type G0 with  $m_V = 10.43$ .

We used a four-channel Chevreton photometer at each site. Three of the channels are employed to monitor the stars (target + comparison 1 + comparison 2), and the fourth channel is devoted to measuring the sky level.

A total of 210 hours of photometric data were obtained during the 21 days of the campaign. The integration time was 1 second. This corresponds to a duty cycle of 40% which is typical for the STEPHI campaigns.

A log of the campaign is given in Table 1. The digits in the data set names (in the first column) indicate the date of the start of the observations and the two last letters indicate the site of




**Fig. 1.** Coverage diagram of the STEPHI XIII campaign. Each rectangular strip represents a night of observation. The labels XL, ZN and SP indicate the three observatories. The total number of hours of photometric data is 210 with an effective coverage of 40%. The times of minimum of the primary eclipses are indicated by E1, E2, E3 and E4.

observation: XL = Xinglong, ZN = Obs. del Teide (Izaña), and SP = San Pedro Mártir. Hereafter, we will use these three abbreviations to represent the three sites. Column 2 gives the UT date at the start of the observations and columns 3 and 4 give the HJD time (Heliocentric Julian Date) at the start and end of the observations. The last two columns show the number of useful measurements and the observation length (in hours). The coverage diagram of the campaign is shown in Fig. 1. The letters in the top of the figure indicate the positions of the observations from the three sites. The rectangular strips represent the duration (in hours) of the observations in relation to 0h UT of the date in the vertical axis. The diagram was made this way to be consistent with Table 1. Note that the observations from each site are aligned inside the same approximate range of time. The times of minimum of the primary eclipses discussed in Sec. 5 are indicated in the figure.

## 3. Data reduction

The data reduction followed similar steps as reported in previous STEPHI campaigns (see, for instance, Hernandez et al. 1998). We start with the inspection of the light curve and, when necessary, elimination of bad points and anomalous parts of the dataset (including the parts where the eclipses are occurring). In the second step we proceed to the calibration using measurements of the sky background simultaneously taken by all the four channels, usually taken at the beginning and at the end of the observations. The third step is the subtraction of the sky background. The measurements of the sky background are subtracted from the light curve of each channel. However, due to technical problems, in this campaign the photometer sky channel could not be used for the sky monitoring on some nights. The fourth

| Data set<br>name | Date        | $T_{begin}$<br>245 0000.+<br>(HJD) | $T_{end}$<br>245 0000.+<br>(HJD) | Number<br>of<br>points | Duration<br>(hours) |
|------------------|-------------|------------------------------------|----------------------------------|------------------------|---------------------|
| IC0607ZN         | 07-Jun-2004 | 3164.51865                         | 3164.70145                       | 18111                  | 5.05                |
| IC0607SP         | 07-Jun-2004 | 3164.78305                         | 3164.94251                       | 11327                  | 3.61                |
| IC0608ZN         | 08-Jun-2004 | 3165.44773                         | 3165.70545                       | 10067                  | 2.97                |
| IC0608SP         | 08-Jun-2004 | 3165.74305                         | 3165.97506                       | 19082                  | 5.37                |
| IC0609XL         | 09-Jun-2004 | 3166.19328                         | 3166.30368                       | 9499                   | 2.65                |
| IC0609ZN         | 09-Jun-2004 | 3166.46177                         | 3166.70003                       | 20586                  | 5.72                |
| IC0609SP         | 09-Jun-2004 | 3166.72632                         | 3166.96818                       | 20407                  | 5.67                |
| IC0610XL         | 10-Jun-2004 | 3167.07915                         | 3167.30217                       | 19267                  | 5.35                |
| IC0610ZN         | 10-Jun-2004 | 3167.43952                         | 3167.69994                       | 22485                  | 6.25                |
| IC0610SP         | 10-Jun-2004 | 3167.72274                         | 3167.97030                       | 16098                  | 4.50                |
| IC0611ZN         | 11-Jun-2004 | 3168.44450                         | 3168.69808                       | 21910                  | 6.09                |
| IC0611SP         | 11-Jun-2004 | 3168.69851                         | 3168.97172                       | 23592                  | 6.56                |
| IC0612XL         | 12-Jun-2004 | 3169.08350                         | 3169.29475                       | 16178                  | 5.07                |
| IC0612ZN         | 12-Jun-2004 | 3169.43251                         | 3169.69744                       | 22884                  | 6.36                |
| IC0612SP         | 12-Jun-2004 | 3169.69386                         | 3169.97170                       | 23435                  | 6.52                |
| IC0613ZN         | 13-Jun-2004 | 3170.43495                         | 3170.70100                       | 22986                  | 6.39                |
| IC0613SP         | 13-Jun-2004 | 3170.69817                         | 3170.97177                       | 23591                  | 6.57                |
| IC0614ZN         | 14-Jun-2004 | 3171.42569                         | 3171.69950                       | 23605                  | 6.57                |
| IC0614SP         | 14-Jun-2004 | 3171.74160                         | 3171.96188                       | 18238                  | 5.07                |
| IC0615ZN         | 15-Jun-2004 | 3172.41819                         | 3172.69828                       | 24191                  | 6.72                |
| IC0615SP         | 15-Jun-2004 | 3172.69699                         | 3172.96155                       | 22845                  | 6.35                |
| IC0616ZN         | 16-Jun-2004 | 3173.41763                         | 3173.69907                       | 24256                  | 6.75                |
| IC0616SP         | 16-Jun-2004 | 3173.69528                         | 3173.95722                       | 22574                  | 6.29                |
| IC0617ZN         | 17-Jun-2004 | 3174.41541                         | 3174.69871                       | 24476                  | 6.80                |
| IC0617SP         | 17-Jun-2004 | 3174.68134                         | 3174.96363                       | 24302                  | 6.77                |
| IC0618ZN         | 18-Jun-2004 | 3175.41786                         | 3175.69765                       | 24172                  | 6.72                |
| IC0618SP         | 18-Jun-2004 | 3175.71757                         | 3175.96352                       | 20072                  | 5.58                |
| IC0619ZN         | 19-Jun-2004 | 3176.42122                         | 3176.69892                       | 23990                  | 6.67                |
| IC0619SP         | 19-Jun-2004 | 3176.68105                         | 3176.96442                       | 24959                  | 6.96                |
| IC0620SP         | 20-Jun-2004 | 3177.67549                         | 3177.96542                       | 24959                  | 6.96                |
| IC0621SP         | 21-Jun-2004 | 3178.68041                         | 3178.96473                       | 22922                  | 6.40                |
| IC0622SP         | 22-Jun-2004 | 3179.67059                         | 3179.96465                       | 24506                  | 6.84                |
| IC06S3SP         | 23-Jun-2004 | 3180.68252                         | 3180.96416                       | 24180                  | 6.76                |
| IC0624SP         | 24-Jun-2004 | 3181.67754                         | 3181.95713                       | 23220                  | 6.46                |
| IC0625SP         | 25-Jun-2004 | 3182.67586                         | 3182.96803                       | 25104                  | 7.01                |
| IC0626SP         | 26-Jun-2004 | 3183.68674                         | 3183.70567                       | 23307                  | 6.52                |
| IC0627SP         | 27-Jun-2004 | 3184.67712                         | 3184.89731                       | 21940                  | 6.14                |
| IC0628SP         | 28-Jun-2004 | 3185.67427                         | 3185.96532                       | 14789                  | 4.34                |

**Table 1.** Log of the observations of the STEPHI XIII campaign. The letters XL, ZN and SP in the data set names indicate the site of observation.

step is the division of the light curve of the variable star (var) by the primary comparison star light curve, var/comp1, and by the secondary comparison star light curve, var/comp2. The purpose of this step is to minimize the effect of changes in the sky transparency on long time scales.

In order to remove low-frequency trends which affect the detection of pulsation modes, the next step is to fit and subtract a polynomial of low-order (order  $\leq 2$ ) from the light curve. The last step is to divide the whole light curve by the average value and subtract 1. The resulting light curve is the relative variation in the magnitude of the star in relation to its mean magnitude. Most of the light curves show maximum variations around 0.05 mag, which corresponds to  $\sim 5\%$  relative to the mean intensity. Finally, all times are converted to heliocentric Julian dates (HJD).

#### 4. Frequency analysis

In Fig. 2 we show the periodograms of the light curves of var/comp1 and var/comp2 for San Pedro Mártir and Tenerife data sets for the range of 0 – 400  $\mu\text{Hz}$  with amplitudes given

in ppt (part per thousand). The respective spectral windows are shown on the right side.

To our surprise, the periodogram of the light curve of the Tenerife data shows the presence of a series of at least 12 harmonics of the frequency of  $1 \text{ cd}^{-1}$  ( $f_d = 11.605 \mu\text{Hz}$ ). This is not the case for the San Pedro Mártir data, although the reduction procedure is the same for the two data sets. We discarded the possibility of these harmonics being an artifact of an electronic problem in one of the channels because they are present in the periodograms of the light curves of var/comp1, var/comp2 and comp2/comp1. The harmonics, perhaps, are resulting of an occasional change in the sky transparency at Izaña during the campaign.

In the periodogram of the SP light curve var/comp1, we can clearly see peaks of pulsation modes with high amplitudes and frequencies within the range 150 – 300  $\mu\text{Hz}$ , completely separated from the bump of peaks of low frequencies ( $f < 150 \mu\text{Hz}$ ). The same peaks appear in the periodogram of the light curve of var/comp2 of the same site, proving that they are related to the target star and not to the comparison star. The difference between the relative heights of the bumps in the two cases can




**Fig. 2.** Periodograms of the one-site light curves of var/comp1 and var/comp2 for the San Pedro Mártir and Tenerife data. In the right side are shown the respective one-site spectral windows.




**Fig. 3.** Periodograms of the Tenerife data obtained with polynomial fitting of order 2 (top) and order 3 (bottom).

be explained by the difference in spectral type: the variable star HD 172189 is an A2 star and the first comparison star (comp1) is of spectral class B9, while the second one (comp2) is a G0 star.

The presence of the harmonics complicates the identification of the peaks of pulsation frequencies in the spectrum. We tried a new modification in the data reduction procedure: we fitted and subtracted a polynomial of order three (instead of order two) from the Tenerife data from each night. This procedure was effective in eliminating all the peaks with high amplitude in the low-frequency region of the periodograms (0-100  $\mu\text{Hz}$ ) and the harmonics of the frequency of the day, without any apparent effect over the peaks with higher amplitude in the pulsation range of frequencies (100 – 300  $\mu\text{Hz}$ ) as can be seen in Fig. 3.

In Fig. 4 we show the spectral window (top) and the periodograms of the whole light curve (ZN + SP + XL) of var/comp1 (middle) and var/comp2 (bottom), where we can see the presence of pulsation modes in both cases. The central peak of the spectral window was arbitrarily positioned to coincide with the position of the highest peak in the periodogram. The FWHM of the central peak is 0.54  $\mu\text{Hz}$  and the two nearest




**Fig. 5.** Prewhitening process in HD 172189 (var/comp1). In each panel the peaks above the 99% confidence level (dashed line) are selected and removed from the original light curve, together with all previous selected frequencies, and a new periodogram is obtained. The selected peaks are indicated by  $f_1$ ,  $f_2$ , etc. The lower level of 40% is indicated by the dotted curve and the solid curve represents the average amplitude.

sidelobes have  $\sim 40\%$  of its amplitude. All subsequent sidelobes have heights below  $\sim 4\%$  of the amplitude of the central peak. The pulsation frequencies appear in both periodograms, in the range 150 – 250  $\mu\text{Hz}$ .

To find pulsation frequencies we used the usual iterative approach: starting with an empty list of candidate frequencies and the periodogram of the original light curve (var/comp1); (1) inside the region of interest in the amplitude spectrum we identify the peaks with the highest confidence levels (taking care to discard aliases). If there is no peak with a significant probability the algorithm stops. (2) Put the selected frequencies in the list of candidate frequencies; (3) using a non-linear method, fit sinusoids with *all* frequencies of the list of frequencies to the original light curve. The fitting refines the values of the initial frequencies and calculates the amplitudes and phases, as well as the respective uncertainties. (4) The fitted sinusoids are subtracted from the original light curve and the periodogram of the residual light curve is calculated, and return to step (1) to search for additional possible pulsation frequencies.

Before applying the algorithm on the light curve var/comp2, we used a high-pass filter to remove signal at low frequencies ( $< 100 \mu\text{Hz}$ ). As in previous STEPPI articles (e.g. Álvarez et al. 1998 Fox-Machado et al. 2002), the confidence levels are calculated with Fisher’s test as prescribed by Nowroozi (1967). We search for oscillation peaks in the range 100 – 600  $\mu\text{Hz}$ , which in Nowroozi’s description corresponds to around  $m = 1000$  “independent” frequency bins. This sets the 99% confidence level at 3.4 times the local square-root of the mean power spectrum  $\sqrt{A_f^2}$ , which is fitted by the exponential  $\bar{A}_f^2 = c_0 \exp(c_1 f)$ , where  $c_0$  and  $c_1$  are constants and  $f$  is the frequency.

The result of the prewhitening process in HD 172189 is shown in Fig. 5. The dashed curve in each graph indicates the 99% confidence level, adopted as a detection limit while the dotted curve indicates the position of the 40% confidence level. In the periodogram of the original light curve, we found three peaks




**Fig. 4.** Spectral window (top) and periodograms of the whole light curves of HD 172189 of var/comp1 (middle) and var/comp2 (bottom).

| $f_i$ | Frequency<br>( $\mu\text{Hz}$ ) | Amplitude<br>(ppt) | Confidence<br>Level |
|-------|---------------------------------|--------------------|---------------------|
| $f_1$ | $226.641 \pm 0.003$             | $2.19 \pm 0.02$    | 99%                 |
| $f_2$ | $200.506 \pm 0.006$             | $0.99 \pm 0.02$    | 99%                 |
| $f_3$ | $206.270 \pm 0.006$             | $0.98 \pm 0.02$    | 99%                 |
| $f_4$ | $196.244 \pm 0.008$             | $0.81 \pm 0.02$    | 99%                 |
| $f_5$ | $215.892 \pm 0.010$             | $0.69 \pm 0.02$    | 99%                 |
| $f_6$ | $225.446 \pm 0.010$             | $0.67 \pm 0.02$    | 99%                 |
| $f_7$ | $242.853 \pm 0.012$             | $0.50 \pm 0.02$    | 65%                 |

**Table 2.** Detected pulsation frequencies in HD 172189 with confidence levels  $\geq 40\%$ .

above the 99% confidence level, indicated in the first graph (top) as  $f_1$ ,  $f_2$ , and  $f_3$ . In the two successive prewhitened periodograms we found three more frequencies above this level,  $f_4$ ,  $f_5$  and  $f_6$ . After the prewhitening of the six frequencies, only one peak was found with a confidence level upper the 40% probability limit, with a 65% confidence level. The six detected pulsation frequencies and the seventh frequency with lower probability are given in Table 2.

We search for the seven detected peaks in the periodogram of the light curve of var/comp2 and using a least square fitting we estimated the probability of each peak not being due to noise in the periodogram of this light curve: 99.9% for  $f_1$ ; 92% for  $f_2$ ; 96% for  $f_3$ ; 80% for  $f_4$ ; 30% for  $f_5$ ; 45% for  $f_6$ ; and 35% for  $f_7$ . As expected, the probabilities found here are lower than those ones found for the case of var/comp1 because the light curve of var/comp2 is more noisy. Even so, the four highest peaks are clearly confirmed in this test.

M-R found two pulsation modes in HD 172189:  $226.82 \mu\text{Hz}$  and  $218.51 \mu\text{Hz}$ . The first mode is in agreement with  $f_1$ , but the second one does not agree with our frequencies. However, we identify the peak at  $218.51 \mu\text{Hz}$  as an alias of  $f_3$ .

## 5. Eclipses in HD 172189

The binarity of HD 172189 was discovered from analysis of the data of a campaign carried out in 1997 for detecting  $\gamma$  Doradus pulsating stars in the cluster IC 4756 (Martín 2000; Martín 2003). M-R analyzed Strömgren  $uvby - H_\beta$  observations obtained in different campaigns during 1997, 2003, and 2004,

showed the presence of three light minima due to eclipses, but only one of the eclipses was followed past the minimum in brightness with a depth of  $\sim 0.12$  mag. From the data analysis, they found a period of  $P = 5.70198$  days. The ephemeris obtained from the data is  $T_{\min I} = 2\,452\,914.644(3) + 5.70198(4) \cdot E$  (in HJD), where  $E$  is the integer number of primary eclipses since the epoch 245 2914.644 HJD. The results indicate a system with an orbital eccentricity of  $e \simeq 0.24$ , a longitude of the periastron of  $\omega \simeq 68^\circ$  and an inclination of  $i \simeq 73^\circ$ . The analysis also suggests that the two stars have different radii ( $r_2/r_1 \simeq 0.6$ ), but similar temperatures  $T_{\text{eff};1} \simeq 1.05 T_{\text{eff};2}$ .

According to the ephemeris, there were four primary eclipses during the present campaign, but, as shown in Fig. 1, only the eclipses  $E1$ ,  $E3$  and  $E4$  occurred while the star was being observed. The first one by the Tenerife observatory and the last two by the Mexican observatory, in the data sets IC0608ZN, IC0619SP and IC0625SP, respectively (cf. Tab 1).

Fig. 6 shows the positions of the eclipses in the light curve of HD 172189. The seven detected pulsation frequencies were subtracted and points collected within 60 s were binned. The three primary eclipses are clearly visible.

Fig. 7 shows the light curves during the eclipse. In the first two panels we see the light decreasing, passing through a minimum in brightness and then increasing. The primary eclipse  $E1$  has its minimum at 2453165.520 (HJD). Unfortunately, the observations were interrupted several times during the night of this eclipse and the reduction of the data presented some difficulties.

A better scenario is shown in the second graph of Fig. 7 for the primary eclipse  $E3$ , where the light curve changes slowly and the minimum is well defined. Fitting a polynomial curve of degree 3 to the eclipse part of the light curve, we obtain  $T_{\min I} = 2453176.920(2)$  (HJD), which differs by  $\sim 9\sigma$  from the predicted value using the ephemeris from M-R, indicating that the uncertainty in the orbital period is underestimated. Using this new minimum and the previous ephemeris, we recalculated the orbital period to obtain  $P = 5.70165(8)$  days. Adopting the instant of the primary eclipse  $E3$  as the new epoch, the new ephemeris is:

$$T_{\min I} = 2453176.920(2) + 5.70165(8) \cdot E \quad (\text{in HJD}). \quad (1)$$

Table 3 shows the ephemeris dates for the four eclipses. The calculated date for the eclipse  $E1$  is 245 3165.518(2), consistent with the minimum observed in the light curve. The maximum in




**Fig. 6.** Light curve of HD 172189. The measurements were binned to 60 seconds. Three primary eclipses, E1, E3 and E4 are clearly visible.




**Fig. 7.** The three primary eclipses (E1, E3 and E4) observed during the campaign. The points were binned and the pulsation frequencies were subtracted.

| Eclipse | Date (HJD)      |
|---------|-----------------|
| E1      | 245 3165.517(2) |
| E2      | 245 3171.218(2) |
| E3      | 245 3176.920(2) |
| E4      | 245 3182.612(2) |

j- ! RIGHT date

**Table 3.** Primary eclipses of HD 172189.

depth for the primary eclipses E1 and E3 are  $\Delta mag = 0.10(3)$  and  $\Delta mag = 0.08(3)$ , respectively, in agreement with the minimum of  $\Delta mag = 0.12$  for the eclipse observed by M-R.

The light curves show that the duration of each primary eclipse is  $\sim$  seven hours and the half-depths occur around one hour before and after the instant of minimum. From the orbital elements calculated by M-R, we calculated a phase of  $\sim 0.132$  for the secondary eclipse, i.e., the secondary eclipses would occur  $\sim 0.75$  days ( $\sim 18$  hours) after each primary eclipse and have the same duration. We searched for minima in the light curve around these dates, but as was remarked by M-R in their analy-

sis, no minimum with an unequivocal and statistically significant depth was found.

## 6. Conclusions

The STEPHI XIII campaign carried out in June 2004, allowed a study of the pulsating behavior of the  $\delta$  Scuti star HD 172189. The Fourier analysis of the light curve from three sites shows the presence of six pulsation frequencies with a confidence level of 99%. An additional frequency with a lower probability of 65% was also found. The observation of three eclipses during the campaign — two of them including the minimum in intensity — allows us to refine the previous value for orbital period of the system, obtaining 5.70165(8) days.

With the detection of six pulsation frequency with a high confidence level, our present campaign confirms the interest of HD 172189 as eclipsing binary featuring a  $\delta$  Scuti pulsator for seismology. Besides the number of modes detected, the interest of HD 172189 Compared with other objects like RZ Cas (Rodríguez et al. 2004a) or AB Cas (Rodríguez et al. 2004b) is also associated with its orbital period. Both, AB Cas and RZ Cas, reveal a short orbital period ( $\sim 1$  d). This suggests possible important effect of tidal forces on the star shape and structure, potential effects on oscillations. In fact, both these objects are considered Algol-type binaries, which supposes that mass transfer occurred between companions and changed their evolution compared with single stars. With a 5.7d orbital period, HD 172189 is expected to be more representative of a "normal" single star, both for its pulsational behavior and for its evolution history.

Dedicated spectroscopic and photometric observations are under way to characterize precisely the orbital parameters and radii of HD 172189 members. These results when available, completed by our present oscillation frequencies analysis will allow to start the modeling and seismic analysis of this object taking advantage of these numerous constraints. In addition to this, within a few years now, it is planed to observe HD 172189 with CoRoT for five months. This is expected to allow detection of many more oscillation modes by pushing down the noise level by a factor 500 in the power spectrum (see Michel et al. 2006). Beside this, the fact that HD 172189 belongs to the cluster IC 4756, even if this one is not so well studied yet, offers an interesting potential source of complementary information for further developments.

**Acknowledgements.** This work received financial support from the Spanish DGES (ESP2004-03855-C03-03), the Chinese National Natural Science Foundation under grant number 10573023 and 10433010, and the Brazilian agency - Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq. The 1.5 m Carlos Sánchez Telescope is operated on the island of Tenerife by the Instituto de Astrofísica de Canarias at the Spanish Observatorio del Teide.

## References

- Alecian, E., Goupil, M.-J., Lebreton, Y., et al. 2006, in Proc. IAU Symp. 240 - Binary Stars as Critical Tools and Tests in Contemporary Astrophysics, 2006, Prague, Czech Republic, p.176
- Álvarez, M., Hernández, M. M., Michel, E., et al. 1998, A&A, 340, 149
- Aerts, C. and Harmanec, P. 2004, in ASP Conf. Ser. 318: Spectroscopically and Spatially Resolving the Components of the Close Binary Stars, 2004, eds. Hilditch, R. W., Hensberge, H. and Pavlovski, K., p. 325
- Aerts, C. 2006, in Proc. IAU Symp. 240 - Binary Stars as Critical Tools and Tests in Contemporary Astrophysics, 2006, Prague, Czech Republic, p.40
- Brown, T.M., Christensen-Dalsgaard, J., Weibel-Mihalas, B., Gilliland, R. L. 1994, ApJ, 427, 1013
- Creevey, O. L., Brown, T. M., Jiménez-Reyes, S., Belmonte, J. A. 2006, in ASP Conference Ser. 349: Astrophysics of Variable Stars, eds. C. Sterken and C. Aerts, p.387

- Fox-Machado, L., Álvarez, M., Michel, E., et al. 2002, A&A, 382, 556-562  
Hernandez, M. M., Michel, E., Belmonte, J. A., et al. 1998, A&A, 337, 198-206  
Herzog, A. D., Sanders, W. L., and Seggewiss, W. 1975, A&AS, 19, 211  
Kopff, E. 1943, Astronomische Nachrichten, 274, 69  
Martín, S. 2000, Ph.D. Thesis, University of Granada, Spain  
Martín, S. 2003, in ASP Conf. Ser. 292: Interplay of Periodic, Cyclic and Stochastic Variability in Selected Areas of the H-R Diagram, eds. C. Sterken, p.59  
Martín-Ruiz, S., Amado, P.J., Suárez, et al. 2005, A&A, 440, 711-714  
Michel, E., Belmonte, J.A., Álvarez, M., et al. 1992, A&A, 255, 139  
Michel, E., Chevreton, M., Belmonte, J. A., et al. 2000, in ASP Conf. Ser. 203: The Impact of Large-Scale Surveys on Pulsating Star Research, eds. L. Szabados and D. Kurt, p.483  
Michel, E. et al. 2006, in Vienna Workshop on the Future of Asteroseismology, Sept 2006, in press  
Miglio, A., Montalbán, J. 2005, A&A, 441, 615  
Nowroozi, A.A. 1967, Geophys. J. R. astr. Soc., 12, 517-720  
Rodríguez, E., García, J. M., Mkrichian, D. E., et al. 2004, MNRAS, 347, 1317  
Rodríguez, E., García, J. M., Gamarova, A. Y., et al. 2004, MNRAS, 353, 310  
Seggewiss, W. 1968, Mitt. Astron. Ges. 25, 122

## List of Objects

- ‘HD 172189’ on page 1  
‘HD 172189’ on page 1  
‘IC 4756’ on page 1  
‘HD 172189’ on page 1  
‘HD 172189’ on page 1  
‘IC 4756 93’ on page 1  
‘SAO 123754’ on page 1  
‘HD 172189’ on page 1  
‘HD 172189’ on page 1  
‘HD 172189’ on page 1  
‘IC 4756’ on page 1  
‘HD 172189’ on page 1  
‘IC 4756’ on page 1  
‘HD 172189’ on page 1  
‘HD 172189’ on page 2  
‘IC 4756’ on page 2  
‘HD 172248’ on page 2  
‘IC 4756 117’ on page 2  
‘SAO 123762’ on page 2  
‘IC 4756 142’ on page 2  
‘IC 4756 115’ on page 2  
‘HD 172189’ on page 4  
‘HD 172189’ on page 5  
‘HD 172189’ on page 5  
‘HD 172189’ on page 5  
‘HD 172189’ on page 5  
‘HD 172189’ on page 6  
‘HD 172189’ on page 6  
‘HD 172189’ on page 6  
‘HD 172189’ on page 6  
‘HD 172189’ on page 6  
‘IC 4756’ on page 6